

Second order linear differential equations with analytic uncertainties: stochastic analysis via the computation of the probability density function

M. Jornet^{a,*}, J. Calatayud^a, O.P. Le Maître^b, J.-C. Cortés^a

^a*Institut Universitari de Matemàtica Multidisciplinar, Universitat Politècnica de València, Camí de Vera s/n, 46022, Valencia, Spain*

^b*CMAP, CNRS, INRIA, École Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau, France*

Abstract

This paper concerns the analysis of random second order linear differential equations. Usually, solving these equations consists of computing the first statistics of the response process, and that task has been an essential goal in the literature. A more ambitious objective is the computation of the solution probability density function. We present advances on these two aspects in the case of general random non-autonomous second order linear differential equations with analytic data processes. The Fröbenius method is employed to obtain the stochastic solution in the form of a mean square convergent power series. We demonstrate that the convergence requires the boundedness of the random input coefficients. Further, the mean square error of the Fröbenius method is proved to decrease exponentially with the number of terms in the series, although not uniformly in time. Regarding the probability density function of the solution at a given time, we rely on the law of total probability to express it in closed-form as an expectation. For the computation of this expectation, a sequence of approximating density functions is constructed by reducing the dimensionality of the problem using the truncated power series of the fundamental set. We prove several theoretical results regarding the pointwise convergence of the sequence of density functions and the convergence in total variation. The pointwise convergence turns out to be exponential under a Lipschitz hypothesis. As the density functions are expressed in terms of expectations, we propose a symbolic Monte Carlo sampling algorithm for their estimation. This algorithm is implemented and applied on several numerical examples designed to illustrate the theoretical findings of the paper.

Keywords: random non-autonomous second order linear differential equation, mean square analytic solution, probability density function, classical Monte Carlo simulation, uncertainty quantification

2010 MSC: 37H10, 34F05, 60H10, 60H35, 65C05

*Corresponding author

Email addresses: marcjornet31@gmail.com (M. Jornet), jucagre@doctor.upv.es (J. Calatayud), olivier.le-maitre@polytechnique.edu (O.P. Le Maître), jccortes@imm.upv.es (J.-C. Cortés)

1. Introduction

Random differential equations are ordinary differential equations whose input coefficients are random quantities, in the form of random variables or stochastic processes (not to be confused with Itô's stochastic differential equations, which are differential equations driven by Wiener noise). In this setting, there is a complete abstract probability space $(\Omega, \mathcal{F}, \mathbb{P})$, where Ω is the sample space defined as the set of outcomes $\omega \in \Omega$, $\mathcal{F} \subseteq 2^\Omega$ is the σ -algebra of events, and \mathbb{P} is the probability measure.

The stochastic solution may be considered in two senses. One approach considers the random calculus that arises from the Lebesgue space $(L^p(\Omega), \|\cdot\|_p)$, $1 \leq p \leq \infty$, where the norms are defined as $\|X\|_p = \mathbb{E}[|X|^p]^{1/p}$ for $p < \infty$, and $\|X\|_\infty = \inf\{C > 0 : |X| \leq C \text{ almost surely}\}$ (essential supremum), where \mathbb{E} denotes the expectation operator defined as $\mathbb{E}[X] = \int_\Omega X \, d\mathbb{P}$. The limits in the definitions of continuity, differentiability, and Riemann integrability are considered in the topology of $L^p(\Omega)$. The space $(L^p(\Omega), \|\cdot\|_p)$, $1 \leq p \leq \infty$, is a Banach space. Of particular importance is the case $p = 2$, which gives rise to the Hilbert space $(L^2(\Omega), \langle \cdot, \cdot \rangle)$ of random variables with finite variance, whose inner product is defined as $\langle X, Y \rangle = \mathbb{E}[XY]$, $X, Y \in L^2(\Omega)$. The calculus in $L^2(\Omega)$ is referred to as the mean square calculus. A key feature of $L^2(\Omega)$ is that mean square convergence ensures convergence of the expectation and the variance. An alternative strategy to tackle random differential equations is the sample path approach, which considers the trajectories of the solution process by fixing each outcome $\omega \in \Omega$. An interesting result that links $L^p(\Omega)$ and sample path calculus states that every $L^p(\Omega)$ solution is also a sample path solution. For theoretical discussions on random differential equations and the types of stochastic solutions, we refer the reader to [1, 2, 3, 4].

Understanding the inherent stochastic nature of the solution is of primary importance. This is the focus of uncertainty quantification [5]. The most common strategies for uncertainty quantification are Monte Carlo simulation [6], PC (polynomial chaos) expansions [7, 8, 9], and perturbation methods [1, 8, 10]. Some studies of different random differential equation problems providing a fair overview of the state-of-the-art literature can be found, for instance, in [11, 12, 13, 14, 15, 16].

In the case of random second order linear differential equations, important advances have been achieved for the computation of the first moments of the solution, via mean square calculus and the so-called Fröbenius method. The Fröbenius method consists in finding a mean square convergent power series solution, in analogy to the deterministic theory of ordinary differential equations. The general stochastic system is given by

$$\begin{cases} \ddot{X}(t) + A(t)\dot{X}(t) + B(t)X(t) = 0, & t \in \mathbb{R}, \\ X(t_0) = Y_0, \quad \dot{X}(t_0) = Y_1. \end{cases} \quad (1)$$

Here, $A(t)$ and $B(t)$ are stochastic processes and Y_0 and Y_1 are random variables on $(\Omega, \mathcal{F}, \mathbb{P})$. The stochastic process $X(t)$ is the solution. We will assume that $A(t)$ and $B(t)$ are analytic stochastic processes on $(t_0 - r, t_0 + r)$, for $r > 0$ fixed, in the mean square sense [1, p. 99]: $A(t) = \sum_{n=0}^{\infty} A_n(t - t_0)^n$ and $B(t) = \sum_{n=0}^{\infty} B_n(t - t_0)^n$ are two random power series in $L^2(\Omega)$, where $A_0, A_1, \dots, B_0, B_1, \dots$ are second order random variables. The expansions coincide with the Taylor series of $A(t)$ and $B(t)$.

Airy, Hermite and Legendre differential equations are particular instances of (1), which represent important stochastic models of Mathematical Physics. The rigorous analysis and construction of mean square solutions to these particular equations, using random power series, can be

found in [17, 18, 19]. We have proposed a generalization of these contributions to the general system (1) in [20, 21].

In the recent paper [22], we investigated the resolution of random second order linear differential equations with PC-based methods. In [23], the authors proposed a homotopy technique to solve some particular random differential equations pertaining to the class given in (1). Other solution techniques include variational iteration [24] and Adomian decomposition [25]. Finally, a technique, analogous to the Fröbenius method but relying on the concept of differential transform, is proposed in [26, 27].

A more ambitious objective is the computation of the probability density function of $X(t)$, denoted hereafter as $f_{X(t)}(x) = \frac{d(\mathbb{P} \circ X(t)^{-1})(x)}{dx}$. The probability density function is defined as a non-negative Borel measurable function characterized by $\mathbb{P}[X(t) \in C] = \int_C f_{X(t)}(x) dx$. Random variables having a probability density function are called absolutely continuous, meaning that their probability law is absolutely continuous with respect to the Lebesgue measure. The density function allows calculating general statistics (expectation, variance, skewness, kurtosis, median, quantiles, mode, etc.) and confidence intervals via integration.

In [28], the authors constructed approximations of the probability density functions of the solution to (1) when $A(t)$ and $B(t)$ do not vary stochastically in time, that is, when $A(t) = A$ and $B(t) = B$ are actually absolutely continuous random variables (autonomous case). A recent paper, [29], presents the approximation of the probability density function of $X(t)$ when $A(t) = p(t; D)$ and $B(t) = q(t; D)$, that is to say, when both $A(t)$ and $B(t)$ depend on a unique absolutely continuous random variable D . This approach does not extend to the general problem (1) and certain theoretical points from that contribution are unclear.

In this work, we provide an analysis of (1) via the Fröbenius method. The solution is expressed in the form of a mean square convergent power series, under $L^\infty(\Omega)$ convergence of $A(t) = \sum_{n=0}^{\infty} A_n(t - t_0)^n$ and $B(t) = \sum_{n=0}^{\infty} B_n(t - t_0)^n$ and mean square integrability of the initial data Y_0 and Y_1 . The boundedness of the coefficients $A_0, A_1, \dots, B_0, B_1, \dots$ is necessary, as shown in examples of the paper. Truncation of unbounded supports of random coefficients can be carried out to assure the required boundedness. The bias error of the Fröbenius method is proved to decrease exponentially with the number of terms in the series. Therefore rapid approximations of the statistical moments of $X(t)$ can be derived. However, the exponential convergence is not uniform in time, and it may deteriorate as we move away from the initial instant t_0 . Section 2 considers these issues. An additional issue is the computation of the probability density function of $X(t)$. Theoretically, it is given by a closed-form expression in terms of an expectation derived from the law of total probability and by exploiting the linearity of the problem. However, to evaluate it in practice, a dimensionality reduction of the problem is required. By truncating the power series, we construct a sequence of probability density functions that, under certain assumptions regarding Nemytskii operators, converges to the target density function pointwise. In this setting, the pointwise convergence of the densities implies convergence in $L^1(\mathbb{R})$ (total variation distance), and in fact, in $L^p(\mathbb{R})$, for $1 \leq p < \infty$. The pointwise convergence rate is proved to be exponential under a certain Lipschitz condition, albeit being again not uniform in time. This theoretical analysis on the approximation of the probability density function is presented in Section 3. As each approximating density function is expressed in terms of an expectation, they can be estimated via a Monte Carlo sampling strategy. The procedure is implemented in the form of a symbolic algorithm, whose computational aspects are detailed in Section 4. The proposed algorithm is tested on several numerical examples in Section 5, to verify the theoretical findings of the paper and to illustrate computational aspects. Finally, Section 6 draws the main

conclusions and points out potential lines of research for the future.

2. Stochastic solution

The initial value problem (1) was previously studied in the mean square sense. We start by recalling the mean square existence and uniqueness theorem proved in our recent contributions [20, 21]. The proof uses fundamental results from deterministic power series extended to the random scenario, together with the basics of difference equations.

Theorem 2.1. [21, Th. 2] *Let $A(t) = \sum_{n=0}^{\infty} A_n(t-t_0)^n$ and $B(t) = \sum_{n=0}^{\infty} B_n(t-t_0)^n$ be two random series in the $L^\infty(\Omega)$ setting, for $t \in (t_0 - r, t_0 + r)$, being $r > 0$ finite and fixed. Assume that the initial conditions Y_0 and Y_1 belong to $L^2(\Omega)$. Then the stochastic process $X(t) = \sum_{n=0}^{\infty} X_n(t-t_0)^n$, $t \in (t_0 - r, t_0 + r)$, where $X_0 = Y_0$, $X_1 = Y_1$ and for $n \geq 0$, $X_{n+2} = \frac{-1}{(n+2)(n+1)} \sum_{m=0}^n [(m+1)A_{n-m}X_{m+1} + B_{n-m}X_m]$, is an analytic solution to the random initial value problem (1) in the mean square sense. Moreover, it is unique. Furthermore, by [20, Subsection 3.4], if Y_0 and Y_1 are bounded random variables, then $X(t)$ is an analytic $L^\infty(\Omega)$ solution to (1).*

From this fundamental theorem we can extend the theory to a more general convergence measure, by considering $L^p(\Omega)$ convergence, $1 \leq p \leq \infty$: if $A(t)$ and $B(t)$ are two random power series with convergence in $L^\infty(\Omega)$, for $t \in (t_0 - r, t_0 + r)$, and the initial conditions Y_0 and Y_1 belong to $L^p(\Omega)$, then the stochastic process $X(t) = \sum_{n=0}^{\infty} X_n(t-t_0)^n$ is the $L^p(\Omega)$ solution to (1) on $(t_0 - r, t_0 + r)$.

Regarding the rapidity of convergence of the power series $X(t) = \sum_{n=0}^{\infty} X_n(t-t_0)^n$ introduced in Theorem 2.1, some theoretical estimates were obtained in [20, Subsection 3.6], although no rate of convergence was derived. Fixed $r > 0$ finite, given $\rho := |t - t_0| < r$ and given an arbitrary s such that $\rho < s < r$, the following estimate holds:

$$\|X^N(t) - X(t)\|_2 \leq K \left(r, s, \{\|A_i\|_\infty\}_{i=1}^\infty, \{\|B_i\|_\infty\}_{i=1}^\infty, \|Y_0\|_2, \|Y_1\|_2 \right) \cdot \frac{(\rho/s)^{N+1}}{1 - \rho/s}.$$

In general, the estimate holds for p -norms. The constant K can be constructed as follows (see [20]):

- Step 1. Given $u = (r+s)/2 \in (s, r)$, choose a constant $C_u > 0$ such that $\|A_i\|_\infty \leq C_u/u^i$ and $\|B_i\|_\infty \leq C_u/u^i$, $i \geq 0$. Such a constant C_u exists because $\sum_{i=0}^{\infty} \|A_i\|_\infty u^i < \infty$ and $\sum_{i=0}^{\infty} \|B_i\|_\infty u^i < \infty$.
- Step 2. Pick an integer $n \geq 0$ such that $\frac{ns}{(n+2)u} + \frac{C_u s}{n+2} + \frac{C_u s^2}{(n+2)(n+1)} < 1$.
- Step 3. Take $K = \max_{0 \leq m \leq n} H_m s^m$, where $\{H_m\}_{m=0}^\infty$ satisfies the recursive equation: $H_0 = \|Y_0\|_2$, $H_1 = \|Y_1\|_2$ and for $m \geq 0$, $H_{m+2} = \left(\frac{m}{(m+2)u} + \frac{C_u}{m+2} \right) H_{m+1} + \frac{C_u}{(m+2)(m+1)} H_m$.

From the constructed K and given a target error $\epsilon > 0$, a truncation order N satisfying $N > \log(\epsilon^{-1} K (1 - \rho/s)^{-1}) / \log(s/\rho) - 1 = \mathcal{O}(\log(\epsilon^{-1}))$ guarantees a mean square error $\|X^N(t) - X(t)\|_2$ less than ϵ . The number s is arbitrary in (ρ, r) . Unfortunately, we are not aware of any method to choose the optimal $s \in (\rho, r)$ minimizing $N = \log(\epsilon^{-1} K (1 - \rho/s)^{-1}) / \log(s/\rho)$.

We stress several new consequences from these estimates. First, the rate of convergence of $\{X^N(t)\}_{N=0}^\infty$ towards $X(t)$ as $N \rightarrow \infty$ is exponential, for $t \in (t_0 - r, t_0 + r)$, because it is proportional to $(\rho/s)^N$. Second, the convergence rate may deteriorate severely for large $\rho = |t - t_0|$ and large norms of the random input coefficients. Indeed, K is growing with $s > \rho$ and the norms.

The fact that the convergence rate deteriorates for large $|t - t_0|$ is clear. Assume that we have $\sum_{n=N}^{\infty} \|X_n\|_2 |t - t_0|^n < \epsilon$, for some target error $\epsilon > 0$. As $|t - t_0|^n$ increases when $|t - t_0|$ grows, a larger N is needed to achieve a mean square error less than ϵ . This fact may especially occur for $|t - t_0| \geq 1$, as in this case $|t - t_0|^n$ does not tend to 0 when $n \rightarrow \infty$, therefore a faster decay of the coefficients $\|X_n\|_2$ is needed to assure convergence.

The numerical experiments presented in [20] also permitted analyzing the behavior of convergence. As theoretically expected by our exposition, the most important phenomenon observed was that the convergence rate deteriorates severely when the distance $|t - t_0|$ increases, therefore making the Fröbenius method computationally intractable. This issue also occurs with PC-type methods, which require large orders for long-time integration [30]. Nonetheless, for not too large $|t - t_0|$ (this “not too large” is problem-dependent), the Fröbenius method works very well.

Having analyzed the convergence rate of the Fröbenius method, let us focus now on the assumptions of Theorem 2.1. In [21], the following open problem was raised: “If there exists a point $t_1 \in (t_0 - r, t_0 + r)$ such that $A(t_1) \notin L^\infty(\Omega)$ or $B(t_1) \notin L^\infty(\Omega)$, then there exist two initial conditions $Y_0, Y_1 \in L^2(\Omega)$ such that (1) has no mean square solution on $(t_0 - r, t_0 + r)$ ”. This problem implies that the hypotheses used in Theorem 2.1 are sharp, in the sense that counterexamples exist if any of them is relaxed. The following two examples of (1) with an unbounded input coefficient have no mean square solution $X(t)$. The arguments to prove that these examples have no solution follow the reasoning of [3, Example, pp. 541–542].

Example 2.2 (Non-existence of mean square solution $X(t)$). Consider the initial value problem (1) with $A(t) = 0$, $B(t) = Z$ and the initial conditions $X(t_0 = 0) = Y_0$ and $\dot{X}(t_0 = 0) = 0$. Let $Z < 0$ be an unbounded random variable (for example, $Z = -U$, where U follows an Exponential, Gamma, Poisson, etc. distribution). Suppose that for any initial condition $Y_0 \in L^2(\Omega)$ there is a mean square solution $X(t)$. By [3, Th. 3(a)], every mean square solution to a random differential equation problem is a sample path solution. More specifically, there exists an equivalent stochastic process, product measurable, whose sample paths solve the deterministic counterpart of the problem almost surely. Therefore $X(t)$ is a sample path solution (we choose the appropriate representative of the equivalence class), with $X(t) = Y_0 \cosh(\sqrt{-Z} t)$ for all $t \in \mathbb{R}$, almost surely. Fix $t \neq 0$. Consider the random variable $T = \cosh(\sqrt{-Z} t)$. Notice that $\|T\|_\infty = \infty$. Consider the operator $\Delta : L^2(\Omega) \rightarrow L^2(\Omega)$, $\Delta(Y) = YT$. This operator is linear and continuous, as a consequence of the closed graph theorem. Hence, there is a constant $C > 0$ such that $\|YT\|_2 \leq C\|Y\|_2$, for all $Y \in L^2(\Omega)$. In fact, this inequality holds for any random variable Y (since, if $Y \notin L^2(\Omega)$, then $\|Y\|_2 = \infty$). Let $Y = T^m$. We have $\|T^{m+1}\|_2 \leq C\|T^m\|_2$, which yields $\|T^m\|_2 \leq C^m$. That is, $\|T\|_{2m} \leq C$. Hence, $\|T\|_\infty = \lim_{m \rightarrow \infty} \|T\|_{2m} \leq C$, but this is a contradiction. Thus, we conclude that there must exist an initial condition $Y_0 \in L^2(\Omega)$ such that the stochastic problem has no mean square solution.

The case in which $Z > 0$ is unbounded (let us suppose that Z is Gamma distributed) may be tackled analogously, although with a subtlety. Proceeding again by contradiction, let us suppose that for any initial condition $Y_0 \in L^2(\Omega)$ there exists a mean square solution $X(t)$. By [3, Th. 3(a)], $X(t) = Y_0 \cos(\sqrt{Z} t)$ for all $t \in \mathbb{R}$, almost surely. In contrast with the previous case, now $\cos(\sqrt{Z} t)$ is bounded. As $X(t)$ is mean square differentiable, its mean square derivative must be given by $\dot{X}(t) = -Y_0 \sqrt{Z} \sin(\sqrt{Z} t)$ [31, p. 536]. Fix $t \neq 0$ and let $T = -\sqrt{Z} \sin(\sqrt{Z} t)$. Now we do have that $\|T\|_\infty = \infty$, so the previous reasoning based on the closed graph theorem can be applied to deduce that there exists an initial condition $Y_0 \in L^2(\Omega)$ such that $\dot{X} \notin L^2(\Omega)$. This is a contradiction.

The general case, in which Z is an unbounded random variable, is easily addressed now (this includes, for instance, the case of Gaussian random variables). If Z is unbounded, then it must

be unbounded on the positive or negative axis. Let us suppose it unbounded on the positive axis (the other case is completely analogous). Take $\tilde{\Omega} \subseteq \Omega$ such that $\mathbb{P}[\tilde{\Omega}] > 0$ and $Z(\omega) > 0$ for each $\omega \in \tilde{\Omega}$. Consider the new probability subspace $(\tilde{\Omega}, \mathcal{F}_{\tilde{\Omega}} = \mathcal{F} \cap 2^{\tilde{\Omega}}, \mathbb{P}_{\tilde{\Omega}} = \mathbb{P}|_{\mathcal{F}_{\tilde{\Omega}}})$. We restate the random differential equation problem on this new probability space, where $Z > 0$ is unbounded. The previous case thus applies. Therefore we are done since every mean square solution on Ω must also be a mean square solution on $\tilde{\Omega}$. This analysis terminates the example.

Example 2.3 (Non-existence of mean square solution $X(t)$). Let us consider now problem (1) with $A(t) = Z$, $B(t) = 0$ and the initial conditions $X(t_0 = 0) = 0$, $\dot{X}(t_0 = 0) = Y_1$. Let Z be any unbounded random variable. Suppose that for any initial condition Y_1 , there exists a mean square solution $X(t)$. Let $Y(t) = \dot{X}(t)$, which satisfies $\dot{Y}(t) + ZY(t) = 0$, $Y(t_0 = 0) = Y_1$. By [3, Th. 3(a)], $Y(t) = Y_1 e^{-Zt}$ for all $t \in \mathbb{R}$, almost surely. Fix $t \neq 0$ and let $T = e^{-Zt}$. The random variable T is unbounded. Hence, the same reasoning from Example 2.2 based on the closed graph theorem applies again. We conclude that there must exist $Y_1 \in L^2(\Omega)$ such that $Y(t) \notin L^2(\Omega)$, which is a contradiction, and we are done with this example.

The boundedness of the random input coefficients is crucial to obtain the Lipschitz condition demanded by the general existence and uniqueness theorem for random differential equations [1, pp. 118–119], [3], [21, Th. 4]. In practice, to satisfy this mandatory boundedness, one may truncate the support to a large but bounded interval.

3. Computation of the probability density function

We now turn to the computation of the probability density function of $X(t)$. Having clarified the conditions for the existence of the solution, we start by rewriting $X(t)$ in an alternative form.

Theorem 3.1. *Let $A(t) = \sum_{n=0}^{\infty} A_n(t - t_0)^n$ and $B(t) = \sum_{n=0}^{\infty} B_n(t - t_0)^n$ be two random series in the $L^\infty(\Omega)$ setting, for $t \in (t_0 - r, t_0 + r)$, being $r > 0$ finite and fixed. Assume that the initial conditions Y_0 and Y_1 belong to $L^2(\Omega)$. Then the mean square analytic solution $X(t)$ can be expressed as $X(t) = Y_0 S_0(t) + Y_1 S_1(t)$, $t \in (t_0 - r, t_0 + r)$, where $S_0(t)$ and $S_1(t)$ are random power series solutions to (1) in $L^\infty(\Omega)$ for the deterministic initial conditions $S_0(t_0) = 1$, $\dot{S}_0(t_0) = 0$, and $S_1(t_0) = 0$, $\dot{S}_1(t_0) = 1$, respectively.*

Notice that we write $X(t)$ as a linear combination of the fundamental set $\{S_0(t), S_1(t)\}$. This expression exploits the linearity of the problem. The processes $S_0(t)$ and $S_1(t)$ are random power series in $L^\infty(\Omega)$, $S_0(t) = \sum_{n=0}^{\infty} S_{0,n}(t - t_0)^n$, $S_1(t) = \sum_{n=0}^{\infty} S_{1,n}(t - t_0)^n$, whose coefficients satisfy a difference equation as in Theorem 2.1; for $S_0(t)$ it comes $S_{0,0} = 1$, $S_{0,1} = 0$, and for $n \geq 0$, $S_{0,n+2} = \frac{-1}{(n+2)(n+1)} \sum_{m=0}^n [(m+1)A_{n-m}S_{0,m+1} + B_{n-m}S_{0,m}]$, while for $S_1(t)$ we have $S_{1,0} = 0$, $S_{1,1} = 1$, and $S_{1,n+2} = \frac{-1}{(n+2)(n+1)} \sum_{m=0}^n [(m+1)A_{n-m}S_{1,m+1} + B_{n-m}S_{1,m}]$, for $n \geq 0$.

The following lemma is necessary to compute the probability density function $f_{X(t)}(x)$. Its proof is a consequence of the law of total probability [32, Ch. 6], [33, Def. 7.11].

Lemma 3.2. *Let U be an absolutely continuous random variable, independent of the random vector (Z_1, Z_2) , where $Z_1 \neq 0$ almost surely. Then $Z_1 U + Z_2$ is absolutely continuous, with density function $f_{Z_1 U + Z_2}(z) = \mathbb{E}[f_U((z - Z_2)/Z_1)/|Z_1|]$.*

This lemma provides an alternative to the random variable transformation method [29, Th. 1], in the case of affine mappings. It does not require that the random quantities have an absolutely continuous probability law, a fact that presents advantages from the practical perspective. The drawback is that we need independence between U and (Z_1, Z_2) to represent the probability density function as an expectation. The expectation can be approximated via sampling-based statistical methods, as discussed later on. The following theorem, which derives the probability density function of $X(t)$, is a straightforward consequence of Lemma 3.2.

Theorem 3.3. *Let $A(t) = \sum_{n=0}^{\infty} A_n(t - t_0)^n$ and $B(t) = \sum_{n=0}^{\infty} B_n(t - t_0)^n$ be two random series in the $L^\infty(\Omega)$ setting, for $t \in (t_0 - r, t_0 + r)$, being $r > 0$ finite and fixed. Suppose that the initial conditions Y_0 and Y_1 belong to $L^2(\Omega)$. If $S_0(t) \neq 0$ almost surely, if Y_0 is absolutely continuous, with density function f_{Y_0} , and it is independent of the rest of random input parameters of (1), then the mean square solution $X(t)$ has for probability density function*

$$f_{X(t)}(x) = \mathbb{E} \left[f_{Y_0} \left(\frac{x - Y_1 S_1(t)}{S_0(t)} \right) \frac{1}{|S_0(t)|} \right]. \quad (2)$$

An important issue with expression (2) is that $S_0(t)$ and $S_1(t)$ are given by infinite series, therefore truncated approximations are needed. We have to justify that it is legitimate to reduce dimensionality and use truncated random power series for $S_0(t)$ and $S_1(t)$. In what follows, we denote by $S_0^N(t)$ and $S_1^N(t)$ the N -th partial sums of $S_0(t)$ and $S_1(t)$, respectively, which converge in $L^\infty(\Omega)$ for each t . Let $X^N(t) = Y_0 S_0^N(t) + Y_1 S_1^N(t)$ be a truncation of the solution $X(t)$, which converges in the mean square sense for each time t .

In the study of random differential equation problems with no closed-form expression of the solution process but only an infinite expansion, one usually constructs an approximating sequence of stochastic processes with reduced dimensionality and computable probability density function. One thus obtains an approximating sequence of probability density functions, which hopefully presents rapid convergence to the target density function. Moreover, the discontinuity and non-differentiability points of the target density function are captured with no difficulty. In the literature, one may find applications of this type of strategy with power series and Karhunen-Loève developments [15, 34], finite difference schemes [12], and PC expansions [35].

If $S_0^N(t) \neq 0$ almost surely, the probability density function of $X^N(t)$ is

$$f_{X^N(t)}(x) = \mathbb{E} \left[f_{Y_0} \left(\frac{x - Y_1 S_1^N(t)}{S_0^N(t)} \right) \frac{1}{|S_0^N(t)|} \right]. \quad (3)$$

This expression involves a maximum of $2N + 3$ random variables ($Y_1, S_{0,n}$ and $S_{1,n}$ for $0 \leq n \leq N$). Thus, the expectation can be computed by numerical integration (in the case of absolutely continuous random input coefficients), or by a Monte Carlo procedure [8, pp. 53–54], by sampling realizations of $Y_1, S_0^N(t)$ and $S_1^N(t)$. This is the same strategy as the one followed in our recent paper [15]. The approach based on numerical integration would be feasible only in the case of small N and $A(t) = p(t; D), B(t) = q(t; D)$ (D random), as in [29], otherwise it is impractical. This is because the integration dimension relies on the dimension of the random space (the total number of input random variables). The Monte Carlo strategy can cope with high uncertainty dimension and large N , albeit at the expense of introducing a statistical error due to sampling, in addition to the bias error. The sampling error is reduced as the number of realizations increases, but at the cost of higher computational burden.

We need to justify that, for each t , $\lim_{N \rightarrow \infty} f_{X^N(t)}(x) = f_{X(t)}(x)$, for $x \in \mathbb{R}$. This is a strong mode of convergence. Indeed, as we are working with density functions, almost everywhere convergence on \mathbb{R} implies convergence in $L^1(\mathbb{R})$: $\|f_{X(t)} - f_{X^N(t)}\|_{L^1(\mathbb{R})} = \int_{\mathbb{R}} |f_{X(t)}(x) - f_{X^N(t)}(x)| dx \rightarrow 0$ as $N \rightarrow \infty$. This is due to Scheffé's lemma [36, p. 55], [37]. Convergence in $L^1(\mathbb{R})$ is also referred to as convergence in total variation [38, p. 41]: $\|(\mathbb{P} \circ (X^N(t))^{-1}) - (\mathbb{P} \circ (X(t))^{-1})\|_{\text{TV}} := \sup\{|\mathbb{P}[X^N(t) \in F] - \mathbb{P}[X(t) \in F]| : F \in \mathcal{F}\} = \frac{1}{2} \|f_{X^N(t)} - f_{X(t)}\|_{L^1(\mathbb{R})}$. It is also equivalent to convergence in terms of the Hellinger distance [39], $H(\mathbb{P} \circ (X^N(t))^{-1}, \mathbb{P} \circ (X(t))^{-1}) := (1/\sqrt{2}) \|f_{X^N(t)}^{1/2} - f_{X(t)}^{1/2}\|_{L^2(\mathbb{R})}$, via the elementary inequalities $H^2 \leq \|\cdot\|_{\text{TV}} \leq \sqrt{2}H$.

In fact, convergence in $L^1(\mathbb{R})$ may be generalized to convergence in $L^p(\mathbb{R})$, for $1 < p < \infty$, by imposing boundedness on \mathbb{R} of f_{Y_0} . Indeed, in this case, taking a constant $C > 0$ such that $|f_{X^N(t)}(x)| \leq C$ and $|f_{X(t)}(x)| \leq C$, for $N \geq 0$, t and $x \in \mathbb{R}$, the mean value theorem leads to $|\|f_{X^N(t)}\|_{L^p(\mathbb{R})}^p - \|f_{X(t)}\|_{L^p(\mathbb{R})}^p| \leq p C^{p-1} \|f_{X^N(t)} - f_{X(t)}\|_{L^1(\mathbb{R})}$, therefore $\|f_{X^N(t)}\|_{L^p(\mathbb{R})} \rightarrow \|f_{X(t)}\|_{L^p(\mathbb{R})}$ as $N \rightarrow \infty$. By Scheffé's lemma, there is convergence in $L^p(\mathbb{R})$: $\|f_{X(t)} - f_{X^N(t)}\|_{L^p(\mathbb{R})} = (\int_{\mathbb{R}} |f_{X(t)}(x) - f_{X^N(t)}(x)|^p dx)^{1/p} \rightarrow 0$ as $N \rightarrow \infty$.

The pointwise convergence is the object of the following important Theorem 3.7. The result is proved in the spirit of our contribution [15], by utilizing the concept of Nemytskii operator [15, Remark 2.6], [40, pp. 15–17], [41, pp. 154–163].

Lemma 3.4. *Let $\{V_N\}_{N=1}^\infty$ be a sequence of random variables that converges to V in $L^2(\Omega)$. If $\mathbb{P}[V \in \mathcal{D}_{f_{Y_0}}] = 0$, where $\mathcal{D}_{f_{Y_0}}$ is the set of discontinuity points of f_{Y_0} , and if $f_{Y_0}(y) \leq \alpha + \beta y^2$, for certain constants $\alpha, \beta \geq 0$, then $f_{Y_0}(V_N) \rightarrow f_{Y_0}(V)$ as $N \rightarrow \infty$ in $L^1(\Omega)$.*

Proof 3.5. *It suffices to prove that, for every subsequence $\{V_{N_k}\}_{k=1}^\infty$, there exists a subsequence $\{V_{N_{k_l}}\}_{l=1}^\infty$ such that $\lim_{l \rightarrow \infty} f_{Y_0}(V_{N_{k_l}}) = f_{Y_0}(V)$ in $L^1(\Omega)$. Fix any subsequence $\{V_{N_k}\}_{k=1}^\infty$. Since $\lim_{k \rightarrow \infty} V_{N_k} = V$ in $L^2(\Omega)$, by [42, Th. 4.9] there exist a subsequence $\{V_{N_{k_l}}\}_{l=1}^\infty$ and a random variable $\bar{V} \in L^2(\Omega)$ such that $\lim_{l \rightarrow \infty} V_{N_{k_l}}(\omega) = V(\omega)$ and $|V_{N_{k_l}}(\omega)| \leq \bar{V}(\omega)$ almost surely. Since $\mathbb{P}[V \in \mathcal{D}_{f_{Y_0}}] = 0$, the continuous mapping theorem [43, p. 7, Th. 2.3] guarantees that $\lim_{l \rightarrow \infty} f_{Y_0}(V_{N_{k_l}}(\omega)) = f_{Y_0}(V(\omega))$ almost surely. As $f_{Y_0}(V_{N_{k_l}}(\omega)) \leq \alpha + \beta(V_{N_{k_l}}(\omega))^2 \leq \alpha + \beta(\bar{V}(\omega))^2 \in L^1(\Omega)$, we can apply the dominated convergence theorem to conclude that the desired limit holds: $\lim_{l \rightarrow \infty} f_{Y_0}(V_{N_{k_l}}) = f_{Y_0}(V)$ in $L^1(\Omega)$.*

Remark 3.6. *As $S_0(t_0) = 1$ and $S_0(t)$ is continuous in $L^\infty(\Omega)$, we can find a neighborhood of t_0 , say $(t_0 - \delta, t_0 + \delta)$ for certain $\delta > 0$, such that $\|S_0(t) - 1\|_\infty < 1/4$ for all $t \in (t_0 - \delta, t_0 + \delta)$. Hence, $S_0(t) > 3/4 > 0$ almost surely, for $t \in (t_0 - \delta, t_0 + \delta)$. Notice that δ might not be equal to ∞ ; for instance, the deterministic function $X(t) = \sin t$ satisfies $\ddot{X}(t) + X(t) = 0$, $X(t_0 = \pi/2) = 1$ and $\dot{X}(t_0 = \pi/2) = 0$.*

For $t \in (t_0 - \delta, t_0 + \delta)$ fixed, there exists an integer $N_t > 0$ such that $\|S_0^N(t) - S_0(t)\|_\infty < 1/4$, for all $N \geq N_t$. Then $\|S_0^N(t) - 1\|_\infty \leq \|S_0^N(t) - S_0(t)\|_\infty + \|S_0(t) - 1\|_\infty < 1/2$. This implies that $S_0^N(t) > 1/2$ almost surely, $N \geq N_t$. From now on, we will work with $t \in (t_0 - \delta, t_0 + \delta)$.

Theorem 3.7. *Suppose the conditions of Theorem 3.3. If f_{Y_0} is continuous on \mathbb{R} and $f_{Y_0}(y) \leq \alpha + \beta y^2$, for certain constants $\alpha, \beta \geq 0$, then $\lim_{N \rightarrow \infty} f_{X^N(t)}(x) = f_{X(t)}(x)$, for each $t \in (t_0 - \delta, t_0 + \delta)$ and for every $x \in \mathbb{R}$.*

Proof 3.8. *Fix $t \in (t_0 - \delta, t_0 + \delta)$ and $x \in \mathbb{R}$. Let*

$$V_N = \frac{x - Y_1 S_1^N(t)}{S_0^N(t)}, \quad V = \frac{x - Y_1 S_1(t)}{S_0(t)} \quad (4)$$

(here we drop the explicit dependencies of V_N and V on t and x). First, notice that $V_N \rightarrow V$ as $N \rightarrow \infty$ in $L^2(\Omega)$, as an easy consequence of the following facts: $S_0^N(t) > 1/2$ almost surely, for all $N \geq N_t$, $S_0^N(t) \rightarrow S_0(t)$ and $S_1^N(t) \rightarrow S_1(t)$ as $N \rightarrow \infty$ in $L^\infty(\Omega)$, and $Y_1 \in L^2(\Omega)$.

The conditions imposed on f_{Y_0} imply that the Nemytskii operator $V \mapsto f_{Y_0}(V)$ is continuous from $L^2(\Omega)$ to $L^1(\Omega)$, by Lemma 3.4. Hence, $\lim_{N \rightarrow \infty} f_{Y_0}(V_N) \rightarrow f_{Y_0}(V)$ in $L^1(\Omega)$. Since $S_0^N(t) > 1/2$ almost surely, for all $N \geq N_t$, and $\lim_{N \rightarrow \infty} S_0^N(t) \rightarrow S_0(t)$ in $L^\infty(\Omega)$, we deduce that $f_{Y_0}(V_N)/S_0^N(t) \rightarrow f_{Y_0}(V)/S_0(t)$ as $N \rightarrow \infty$ in $L^1(\Omega)$.

In particular, the sequence of expectations, $f_{X^N(t)}(x) = \mathbb{E}[f_{Y_0}(V_N)/S_0^N(t)]$, converges to the density $f_{X(t)}(x) = \mathbb{E}[f_{Y_0}(V)/S_0(t)]$, which completes the proof.

In Section 5, the application of Theorem 3.7 will be illustrated numerically on Examples 5.1–5.2.

Remark 3.9. Having $\lim_{N \rightarrow \infty} f_{Y_0}(V_N)/S_0^N(t) = f_{Y_0}(V)/S_0(t)$ in $L^1(\Omega)$ assures the convergence of the expectations. If convergence of the variances is also needed, one needs to extend the convergence to $L^2(\Omega)$. In this case, the boundedness condition on f_{Y_0} should be $f_{Y_0}(y) \leq \alpha + \beta|y|$ (apply an analogous proof to Lemma 3.4).

Remark 3.10 (Rate of convergence of the approximating density functions). Notice that, under the conditions of Theorem 3.3, if f_{Y_0} is Lipschitz continuous on \mathbb{R} (this assumption is stronger than the hypotheses of Theorem 3.7), then $f_{X^N(t)}(x)$ converges with N exponentially to $f_{X(t)}(x)$, for $t \in (t_0 - \delta, t_0 + \delta)$ and $x \in \mathbb{R}$. This is because the Lipschitz condition allows estimating $|f_{X^N(t)}(x) - f_{X(t)}(x)|$ via the following inequality:

$$|f_{X^N(t)}(x) - f_{X(t)}(x)| \leq C_t \left((|x| + 1) \|S_0^N(t) - S_0(t)\|_\infty + \|Y_1\|_2 \|S_1^N(t) - S_1(t)\|_\infty \right),$$

and as discussed in Section 2, the Fröbenius method converges exponentially. In the previous expression, C_t is a constant depending on t . Unfortunately, the exponential convergence rate is not uniform with t and x . As $|t - t_0|$ grows, one needs to increase N to maintain the accuracy. The same occurs with $|x|$, which increases the bias error $\|S_0^N(t) - S_0(t)\|_\infty$ linearly.

In general, if f_{Y_0} is γ -Hölder continuous on \mathbb{R} with exponent $0 < \gamma \leq 1$ (the case $\gamma = 1$ corresponds to Lipschitz continuity), then

$$|f_{X^N(t)}(x) - f_{X(t)}(x)| \leq C_t \left\{ \|S_0^N(t) - S_0(t)\|_\infty + \left(|x| \|S_0^N(t) - S_0(t)\|_\infty + \|Y_1\|_2 \|S_1^N(t) - S_1(t)\|_\infty \right)^\gamma \right\}.$$

The same conclusion on the convergence holds in this case.

Notice that the regularity of $f_{X^N(t)}(x)$ is inherited from $f_{Y_0}(y)$. These ideas are formalized in the following theorem:

Theorem 3.11. Under the assumptions of Theorem 3.7, if f_{Y_0} is $C^1(\mathbb{R})$ with bounded derivative on \mathbb{R} , then $f_{X^N(t)}(x)$ and $f_{X(t)}(x)$ are $C^1(\mathbb{R})$, with bounded derivatives, and $f'_{X^N(t)}(x) \rightarrow f'_{X(t)}(x)$ as $N \rightarrow \infty$, for each $t \in (t_0 - \delta, t_0 + \delta)$ and for every $x \in \mathbb{R}$.

Proof 3.12. Fix $t \in (t_0 - \delta, t_0 + \delta)$. The following facts permit differentiating under the expectation operator that defines $f_{X^N(t)}(x)$ and $f_{X(t)}(x)$ [44, p. 142]: f_{Y_0} is differentiable with bounded derivative, and $S_0^N(t) > 1/2$ almost surely for all $N \geq N_t$. Hence,

$$f'_{X^N(t)}(x) = \mathbb{E} \left[f'_{Y_0} \left(\frac{x - Y_1 S_1^N(t)}{S_0^N(t)} \right) \frac{1}{(S_0^N(t))^2} \right], \quad f'_{X(t)}(x) = \mathbb{E} \left[f'_{Y_0} \left(\frac{x - Y_1 S_1(t)}{S_0(t)} \right) \frac{1}{(S_0(t))^2} \right].$$

The continuity and boundedness conditions imposed on f'_{Y_0} entail that the Nemytskii operator $V \mapsto f'_{Y_0}(V)$ is continuous from $L^2(\Omega)$ to $L^1(\Omega)$, by Lemma 3.4. Thereby, as in the proof of Theorem 3.7, we deduce that $\lim_{N \rightarrow \infty} f'_{X^N(t)}(x) = f'_{X(t)}(x)$, $x \in \mathbb{R}$.

Remark 3.13. It is important to realize that the previous theory works exchanging the role of Y_1 and Y_0 . Indeed, even though $S_1(t_0) = 0$, in contrast with $S_0(t_0) = 1$, we do have that $\dot{S}_1(t_0) = 1$. We may choose a neighborhood of t_0 , say $(t_0 - \mu, t_0 + \mu)$ for certain $\mu > 0$, such that $\dot{S}_1(t) > 3/4$ almost surely, for $t \in (t_0 - \mu, t_0 + \mu)$. We know that, in the sense of $L^\infty(\Omega)$, $S_1(t) = \int_{t_0}^t \dot{S}_1(r) dr$. Then $|S_1(t)| > \frac{3}{4}|t - t_0| = m_t$ almost surely, for $t \in (t_0 - \mu, t_0 + \mu)$. In particular, as $m_t > 0$ for $t \in (t_0 - \mu, t_0 + \mu) \setminus \{t_0\}$, the previous proofs work with Y_1 in lieu of Y_0 . The previous theoretical results may be restated in a completely analogous fashion, as

$$f_{X(t)}(x) = \mathbb{E} \left[f_{Y_1} \left(\frac{x - Y_0 S_0(t)}{S_1(t)} \right) \frac{1}{|S_1(t)|} \right], \quad f_{X^N(t)}(x) = \mathbb{E} \left[f_{Y_1} \left(\frac{x - Y_0 S_0^N(t)}{S_1^N(t)} \right) \frac{1}{|S_1^N(t)|} \right],$$

for $t \in (t_0 - \mu, t_0 + \mu) \setminus \{t_0\}$. In this case, one requires Y_1 to have an absolutely continuous probability law, with density function f_{Y_1} , and to be independent of the rest of the random input parameters in (1). For convergence, one imposes continuity for f_{Y_1} on \mathbb{R} and boundedness $f_{Y_1}(y) \leq \alpha + \beta y^2$, for certain constants $\alpha, \beta \geq 0$. If f_{Y_1} is Lipschitz continuous on \mathbb{R} , then an exponential convergence holds. Finally, if f_{Y_1} is also $C^1(\mathbb{R})$ with bounded derivative on \mathbb{R} , then both $f_{X^N(t)}(x)$ and $f_{X(t)}(x)$ are $C^1(\mathbb{R})$, with bounded derivative, and the sequence of derivatives converges. These cases are considered in Example 5.3.

The continuity condition on \mathbb{R} imposed in Theorem 3.7 is somewhat restrictive, as we do not allow some common probability distributions for Y_0 whose density function possesses discontinuity points, such as the Uniform, Exponential or general truncated distributions. Notice that this assumption in Theorem 3.7 may be relaxed to almost everywhere continuity on \mathbb{R} , by adding absolute continuity on Y_1 . This fact is a consequence of the continuous mapping theorem [43, p. 7, Th. 2.3]. Indeed, for $t \in (t_0 - \min\{\delta, \mu\}, t_0 + \min\{\delta, \mu\}) \setminus \{t_0\}$, as $|S_1(t)| > m_t > 0$ almost surely and Y_1 is absolutely continuous, then $V = \frac{x - Y_1 S_1(t)}{S_0(t)}$ is absolutely continuous, by Lemma 3.2. Therefore, the probability that V lies in the discontinuity set of f_{Y_0} is 0. This assures that $f_{Y_0}(V_N) \rightarrow f_{Y_0}(V)$ in $L^1(\Omega)$ as $N \rightarrow \infty$, by Lemma 3.4.

The precise restatement of Theorem 3.7 is the following:

Theorem 3.14. Suppose the conditions of Theorem 3.3. If f_{Y_0} is almost everywhere continuous on \mathbb{R} , $f_{Y_0}(y) \leq \alpha + \beta y^2$ for certain constants $\alpha, \beta \geq 0$, Y_1 is absolutely continuous, and Y_1 is independent of (A, B) , then $\lim_{N \rightarrow \infty} f_{X^N(t)}(x) = f_{X(t)}(x)$, for $t \in (t_0 - \min\{\delta, \mu\}, t_0 + \min\{\delta, \mu\}) \setminus \{t_0\}$ and for every $x \in \mathbb{R}$.

Theorem 3.14 will be applied in Example 5.4. An alternative version, with Y_1 playing the role of Y_0 , can be formulated following Remark 3.13. Notice that, nowhere in our theoretical exposition, we require independence between the coefficients of $A(t)$ and $B(t)$. We do not need any assumption on their probability distributions either, which might be discrete or continuous (but always bounded).

The methodology and theory presented in the paper do not cover all situations. For instance, let us study (1) involving discrete uncertainties. Other situations could be analogously analyzed.

Theorem 3.15. *Suppose the conditions of Theorem 3.3. Assume that the coefficients $A_0, A_1, \dots, B_0, B_1, \dots$ are deterministic constants. If f_{Y_0} has at most a countable number of discontinuities on \mathbb{R} , $f_{Y_0}(y) \leq \alpha + \beta y^2$ for certain constants $\alpha, \beta \geq 0$, and Y_1 is a discrete random variable, then $\lim_{N \rightarrow \infty} f_{X^N(t)}(x) = f_{X(t)}(x)$ for almost every $x \in \mathbb{R}$, for each $t \in (t_0 - \delta, t_0 + \delta)$.*

Proof 3.16. *Fix $t \in (t_0 - \delta, t_0 + \delta)$. Let $V_N(x) = (x - Y_1 S_1^N(t))/S_0^N(t)$, $V(x) = (x - Y_1 S_1(t))/S_0(t)$ (now we make the dependence of V_N and V on x explicit). We know that $V_N(x) \rightarrow V(x)$ in $L^2(\Omega)$ as $N \rightarrow \infty$, for all $x \in \mathbb{R}$. Given the discontinuity set of f_{Y_0} , $\mathcal{D}_{f_{Y_0}}$, we need to justify that $\mathbb{P}[V(x) \in \mathcal{D}_{f_{Y_0}}] = 0$, for almost every $x \in \mathbb{R}$. In this case, $f_{Y_0}(V_N(x)) \rightarrow f_{Y_0}(V(x))$ in $L^1(\Omega)$ as $N \rightarrow \infty$, for almost every $x \in \mathbb{R}$.*

Write $\mathcal{D}_{f_{Y_0}} = \{d_1, d_2, d_3, \dots\}$. As Y_1 is a discrete random variable, its support may be expressed as $\mathcal{S}_{Y_1} = \{y_1^1, y_1^2, y_1^3, \dots\}$. Then the support of $V(x)$ is $\mathcal{S}_{V(x)} = \{(x - y_1^j S_1(t))/S_0(t) : j = 1, 2, 3, \dots\}$. The problematic x 's are those such that $x = y_1^j S_1(t) + d_k S_0(t)$. Let $\Lambda = \{y_1^j S_1(t) + d_k S_0(t) : j, k = 1, 2, 3, \dots\}$, which is a countable set. For every $x \notin \Lambda$, $\mathbb{P}[V(x) \in \mathcal{D}_{f_{Y_0}}] = 0$. As a consequence, $\lim_{N \rightarrow \infty} f_{Y_0}(V_N(x)) = f_{Y_0}(V(x))$ in $L^1(\Omega)$, $x \notin \Lambda$, by Lemma 3.4. This gives $\lim_{N \rightarrow \infty} f_{X^N(t)}(x) = f_{X(t)}(x)$, $x \notin \Lambda$, and we are done.

Once again, one can state a similar version with Y_1 playing the role of Y_0 (see Remark 3.13) and working on $(t_0 - \mu, t_0 + \mu) \setminus \{t_0\}$, instead. Example 5.5 covers this situation.

4. Computational aspects

We recast the proposed methodology in the form of Algorithm 1, which corresponds to the case of Y_0 having a density, see (2); following Remark 3.13, one can exchange the role of Y_0 and Y_1 in Algorithm 1, provided that Y_1 has a density.

By judiciously exploiting its expression in (3), $f_{X^N(t)}(x)$ can be approximated *via* a Monte Carlo procedure [8, pp. 53–54] to evaluate the expectation: using M randomly generated realizations of Y_1 , $S_0^N(t)$ and $S_1^N(t)$, we compute the sample average of $V_N(x, t)$ in (4). Algorithm 1 corresponds to symbolic computations with symbolic variables x and t [45]; it computes a function $f_X^{N,M}(x, t)$, which is a complex closed-form expression approximating $f_{X(t)}(x)$. To speed up the execution of the algorithm, numerical values of t and/or x may be provided.

The estimation error can be split into two contributions: $f_{X(t)}(x) - f_X^{N,M}(x, t) = \theta_N(x, t) + \mathcal{E}_{N,M}(x, t)$. The first contribution, $\theta_N(x, t) = f_{X(t)}(x) - f_{X^N(t)}(x)$, is the bias error caused by the truncation order N in the Fröbenius method. It is deterministic and decays exponentially as $N \rightarrow \infty$ for each t and x by Remark 3.10. The second contribution is the sampling error $\mathcal{E}_{N,M}(x, t) = f_{X^N(t)}(x) - f_X^{N,M}(x, t)$, due to using a finite number M of samples (statistical error). This contribution is random and $\mathcal{E}_{N,M}(x, t) \rightarrow 0$ with M almost surely, as a consequence of the law of large numbers. If the variance

$$\sigma_N^2(x, t) := \mathbb{V} \left[f_{Y_0} \left(\frac{x - Y_1 S_1^N(t)}{S_0^N(t)} \right) \frac{1}{S_0^N(t)} \right] \quad (5)$$

is finite, then the asymptotic probability distribution of $\mathcal{E}_{N,M}(x, t)$ as $M \rightarrow \infty$ is, by the central limit theorem, Normal(0, $\sigma_N^2(x, t)/M$). The variance $\sigma_N^2(x, t)$ tends, as $N \rightarrow \infty$, to $\sigma^2(x, t) := \mathbb{V}[f_{Y_0}(\frac{x - Y_1 S_1(t)}{S_0(t)}) \frac{1}{S_0(t)}]$ (see Remark 3.9). In this case, we say that the sampling error is of order $1/\sqrt{M}$, and write $\mathcal{O}(1/\sqrt{M})$. On the contrary, if $\sigma_N^2(x, t) = \infty$, then the almost sure convergence $\mathcal{E}_{N,M}(x, t) \rightarrow 0$ with M remains valid, although it might be much slower and affect the

Algorithm 1 Estimation of $f_{X^N(t)}(x)$ via a classical Monte Carlo procedure.

Inputs: $t_0; N; f_{Y_0}$; probability distribution of $A_0, \dots, A_N, B_0, \dots, B_N, Y_1$; and number M of realizations in the classical Monte Carlo procedure. Here, t and x will be symbolic variables.

Required: Hypotheses of Theorem 3.7 or Theorem 3.14.

- 1: $S_{0,0} \leftarrow 1, S_{0,1} \leftarrow 0, S_{1,0} \leftarrow 0, S_{1,1} \leftarrow 1$ ▷ Initial conditions
- 2: $\Sigma \leftarrow 0$ ▷ Initialize the samples sum
- 3: **for** $i = 1, \dots, M$ **do** ▷ Monte Carlo loop
- 4: Draw randomly a realization of $(A_0, \dots, A_{N-2}, B_0, \dots, B_{N-2})$ and Y_1
- 5: **for** $n = 0, \dots, N - 2$ **do**
- 6: $S_{0,n+2} \leftarrow \frac{-1}{(n+2)(n+1)} \sum_{m=0}^n [(m+1)A_{n-m}S_{0,m+1} + B_{n-m}S_{0,m}]$
- 7: $S_{1,n+2} \leftarrow \frac{-1}{(n+2)(n+1)} \sum_{m=0}^n [(m+1)A_{n-m}S_{1,m+1} + B_{n-m}S_{1,m}]$
- 8: **end for**
- 9: $S_0^N(t) \leftarrow 1 + \sum_{n=1}^N S_{0,n}(t - t_0)^n$ ▷ Realization of $S_0^N(t)$
- 10: $S_1^N(t) \leftarrow \sum_{n=1}^N S_{1,n}(t - t_0)^n$ ▷ Realization of $S_1^N(t)$
- 11: $\Sigma \leftarrow \Sigma + f_{Y_0} \left(\frac{x - Y_1 S_1^N(t)}{S_0^N(t)} \right) \frac{1}{|S_0^N(t)|}$ ▷ Update the samples sum
- 12: **end for**
- 13: $f_X^{N,M}(x, t) \leftarrow \Sigma / M$ ▷ Set sample average
- 14: **Return** $f_X^{N,M}(x, t)$ ▷ Approximation of $f_{X^N(t)}(x)$

approximation to $f_{X^N(t)}(x)$ severely. See the forthcoming Example 5.3 for an illustration of this issue.

Even though the bias error decays very fast, the sampling error is inevitable. In numerical computations, for fixed M , there is usually an index N from which the global error does not go down anymore because the sampling error $O(1/\sqrt{M})$ becomes dominant.

Within the main loop of Algorithm 1 (loop over the samples), we first generate one realization for each random variable $A_0, \dots, A_{N-2}, B_0, \dots, B_{N-2}$ and Y_1 ; these realizations are used to compute by recursion the corresponding realizations of $S_0^N(t)$ and $S_1^N(t)$. In our implementation, this procedure is more effective than expressing first $S_0^N(t)$ and $S_1^N(t)$ recursively in terms of symbolic variables $A_0, \dots, A_{N-2}, B_0, \dots, B_{N-2}$ and Y_1 , and then evaluate for the realizations of $A_0, \dots, A_{N-2}, B_0, \dots, B_{N-2}$ and Y_1 . This is due to the excessive complexity of the symbolic expressions of $S_0^N(t)$ and $S_1^N(t)$, which makes the computational time of their evaluation for specific realizations prohibitively large.

The computational complexity of Algorithm 1 is at most $O(MN^2)$ (the nested loop over n demands $\sum_{n=0}^{N-2} O(n) = O(N^2)$ operations in general). As we show in the following Section 5, the implemented algorithm is certainly applicable and suitable for stochastic computations.

By taking $M = O(1/\epsilon^2)$, the variance of the statistical error is $\mathbb{V}[\mathcal{E}_{N,M}(x, t)] = O(\epsilon^2)$ (assuming the variance in (5) finite). Under exponential convergence of the bias, by picking $N = O(\log(1/\epsilon)) + O(1)$ the bias error is $|\theta_N(x, t)| = O(\epsilon)$. Then the mean square error of the algorithm is $\|f_{X(t)}(x) - f_X^{N,M}(x, t)\|_2 = \sqrt{\theta_N(x, t)^2 + \mathbb{V}[\mathcal{E}_{N,M}(x, t)]} = O(\epsilon)$, with a computational complexity $O(MN^2) = O(\epsilon^{-2} \log^2 \epsilon)$.

The complexity of Algorithm 1 is significantly reduced if $A(t)$ and $B(t)$ are random polynomials, instead of infinite series. Suppose for instance that $A_j = 0$ and $B_j = 0$, for $j \geq N_0 - 1$. Then, within the nested loop over n , we actually sum N_0 terms, instead of n terms. Therefore, the nested loop demands $N_0 O(N) = O(N)$ operations. The whole algorithm then requires $O(MN)$

operations only. If we take $M = O(1/\epsilon^2)$ and $N = O(\log(1/\epsilon)) + O(1)$ to ensure a mean square error of order ϵ , the computational complexity becomes $O(MN) = O(\epsilon^{-2} \log(\epsilon^{-1}))$.

In the case in which $A(t)$ and $B(t)$ are deterministic expansions, the loop over n and the assignments for $S_0^N(t)$ and $S_1^N(t)$ may be run once for all at the beginning of the algorithm and before the loop over the samples. The computational complexity then reduces even more to $O(M) + O(N^2)$ operations and the global cost is generally dominated by the sampling. To guarantee a global mean square error of order ϵ with $M = O(1/\epsilon^2)$ and $N = O(\log(1/\epsilon)) + O(1)$, the computational complexity becomes $O(\epsilon^{-2}) + O(\log^2 \epsilon) = O(\epsilon^{-2})$. This scenario allows increasing M and obtaining more accurate results by improving the statistical convergence. If $A(t)$ and $B(t)$ are simply deterministic polynomials, then the overall cost reduces further to $O(M) + O(N)$ operations, which yields in the end $O(\epsilon^{-2})$ calculations.

In the view of computational applications, an important drawback of our exposition is the lack of awareness on the specific values of δ and μ , which are necessary to prove the theoretical convergence. Given any t , one can apply Algorithm 1 and check the convergence of the estimator with M and N . The results can be validated using other stochastic methods and using statistics based on the estimated density. Notice that, in Algorithm 1, we have put $|S_0^N(t)|$ instead of $S_0^N(t)$. Even though we assume that $S_0^N(t) > 0$ almost surely, for $t \in (t_0 - \delta, t_0 + \delta)$ and $N \geq N_r$, the absolute value ensures positiveness in numerical applications even if $|t - t_0| \geq \delta$.

5. Numerical examples

In this section, we numerically illustrate our theoretical findings, using Algorithm 1 to estimate the density of the solution to (1). Several cases, differing by the probability distributions of the random input coefficients, are considered to cover a large class of situations and show the broad applicability of our theory.

In each of these examples, we first check that the necessary theoretical conditions hold; we then estimate the density function $f_{X^{N(t)}}(x)$ for several increasing values of N to highlight the convergence toward $f_{X(t)}(x)$. To this end, we employ the symbolic Monte Carlo sampling procedure outlined in Algorithm 1.

The theoretical results of Section 3 motivate the structure and the choice of the following five examples. In Example 5.1, we address the case where $A(t)$ and $B(t)$ are random polynomials; while Example 5.2 concerns infinite expansions and infinite dimensionality. These first two examples showcase the applicability of Theorem 3.7. Example 5.3 is designed to highlight Remark 3.13. Up to this example, f_{Y_0} or f_{Y_1} are continuous on the whole real line. In contrast, Example 5.4 considers experiments with f_{Y_0} possessing discontinuity points, thus evoking Theorem 3.14. Finally, Example 5.5 considers the case where $A(t)$ and $B(t)$ are deterministic, so that Theorem 3.15 applies.

The implementations and computations are performed with Mathematica[®], version 11.2 [46], owing to its capability to handle symbolic computations. In general, Algorithm 1 is applied with $M = 20,000$ samples, as beyond this limit, the computational burden is becoming massive. The output function $f_X^{N,M=20,000}(x, t)$ is handled symbolically on t and x . To simplify the notations, we refer to the Monte Carlo estimate $f_X^{N,M=20,000}(x, t)$ as $\hat{f}_{X^{N(t)}}(x)$. We recall that the estimate $\hat{f}_{X^{N(t)}}(x)$ has two sources of error: bias and sampling. Although the bias error decays very fast (exponentially under the conditions of Remark 3.10), the sampling error is unavoidable and at least of order $O(1/\sqrt{M})$.

In each one of the following examples, we perform a complete analysis of the errors. As the exact density function $f_{X(t)}(x)$ is not known, we first analyze differences in consecutive (in N) estimates, both pointwise, using

$$\delta\epsilon^N(x, t) := |\hat{f}_{X^{N+1}(t)}(x) - \hat{f}_{X^N(t)}(x)|, \quad (6)$$

and globally, using the norm

$$\Delta\epsilon^N(t) := \|\hat{f}_{X^{N+1}(t)} - \hat{f}_{X^N(t)}\|_{L^1(\mathbb{R})}. \quad (7)$$

As successive differences do not directly characterize the error, we also report

$$E^N(t) := \|\hat{f}_{X^L(t)} - \hat{f}_{X^N(t)}\|_{L^1(\mathbb{R})} \quad (8)$$

for some pre-fixed $L \gg 1$, selected such that $\hat{f}_{X^L(t)}$ plays the role of a bias-free estimate of function $f_{X(t)}$. We set $L = 30$ in the following. The $L^1(\mathbb{R})$ norms are computed by direct numerical integration, using a standard quadrature rule (standard `NIntegrate` routine in Mathematica[®]).

Example 5.1. We start with the stochastic problem (1) where both $A(t)$ and $B(t)$ are random polynomials of degree 1: $A(t) = A_0 + A_1 t$, and $B(t) = B_0 + B_1 t$. We set $A_0 = 4$, $A_1 \sim \text{Uniform}(0, 1)$, $B_0 \sim \text{Gamma}(2, 2)$, $B_1 \sim \text{Bernoulli}(0.35)$, $Y_0 \sim \text{Normal}(2, 1)$ and $Y_1 \sim \text{Poisson}(2)$, all being independent random variables. In order for the hypotheses of Theorem 2.1 and Theorem 3.7 to be satisfied, the Gamma distribution is truncated. For the Gamma distribution with shape and rate 2, it can be checked that the interval $[0, 4]$ contains approximately 99.7% of the probability, so we actually consider $B_0 \sim \text{Gamma}(2, 2)|_{[0,4]}$.

By Theorem 2.1, the unique mean square solution to the problem can be written as a random power series $X(t) = \sum_{n=0}^{\infty} X_n t^n$ that is mean square convergent for all $t \in \mathbb{R}$. With Theorem 3.7, we approximate pointwise the probability density function $f_{X(t)}(x)$ with $\hat{f}_{X^N(t)}(x)$, $N \geq 0$, and use Algorithm 1 taking advantage from the fact that $A(t)$ and $B(t)$ are random polynomials and not infinite expansions. We consider times $t = 0.5$, 1 and 1.5. In Figure 1 we present the graphs of $\hat{f}_{X^N(t)}(x)$ at the corresponding times. Observe that the estimates are smooth, due to the regularity of the initial density f_{Y_0} , see Theorem 3.11. Observe also that, as N increases, the density functions become closer, reflecting the theoretical convergence. The convergence is made clear in the corresponding successive differences $\delta\epsilon^N(x, t)$ (see (6)) reported in Figure 2. Table 1 presents the $L^1(\mathbb{R})$ norms of the successive differences, $\Delta\epsilon^N(t)$ (see (7)).




Figure 1: Graphical representations of the Monte Carlo estimates $\hat{f}_{X^N(t)}(x)$ at $t = 0.5$ (left), $t = 1$ (center) and $t = 1.5$ (right), with orders of truncation $N = 1-6$, $N = 6-11$ and $N = 11-16$, respectively. This figure corresponds to Example 5.1.




Figure 2: Differences in consecutive estimates $\delta\epsilon^N(x, t)$ (see (6)) at $t = 0.5$ (left), $t = 1$ (center) and $t = 1.5$ (right), with orders of truncation $N = 1-5$, $N = 6-10$ and $N = 11-15$, respectively. This figure corresponds to Example 5.1.

$t = 0.5$	$N = 1$	$N = 2$	$N = 3$	$N = 4$	$N = 5$
	0.903091	0.622968	0.270690	0.0923362	0.0178834
$t = 1$	$N = 6$	$N = 7$	$N = 8$	$N = 9$	$N = 10$
	0.691809	0.263246	0.0912177	0.0345686	0.026688
$t = 1.5$	$N = 11$	$N = 12$	$N = 13$	$N = 14$	$N = 15$
	0.348643	0.180075	0.0721679	0.0320314	0.0198364

Table 1: Norm $\Delta\epsilon^N(t)$ of differences in consecutive estimates (see (7)) for different times t and truncation orders N . This table corresponds to Example 5.1.

The left plot in Figure 3 reports (in log-scale) the error estimate $E^N(t)$ (see (8)), for $t = 0.5$, $t = 1$ and $t = 1.5$. From the plot, it is clear that there is an index N from which the error does not go down anymore, because of the sampling error (recall that we used a fixed number of samples $M = 20,000$). Notice also that, as $|t - t_0| = |t|$ gets larger, a higher order of truncation N is required to enhance the approximations of $f_{X(t)}(x)$. In the right plot of Figure 3, we report the error estimate, $E^N(t)$, as a function of the consecutive difference, $\Delta\epsilon^N(t)$, for $t = 0.5$, $t = 1$ and $t = 1.5$. We also plot a regression line through the data to reflect the exponential relationship between $E^N(t)$ and $\Delta\epsilon^N(t)$, at a given t ,

$$\log E^N(t) \approx \log \beta(t) + \alpha(t) \log \Delta\epsilon^N(t). \quad (9)$$

There are three regression lines, one for each time t . We observe a strong linear relation with N between the errors and the successive differences in log-scale, with slope $\alpha(t)$ being approximately 1, at least up to the truncation order at which the sampling error becomes dominant. This finding suggests that it is possible to estimate the norm of the bias error, $\|\theta_N(\cdot, t)\|_{L^1(\mathbb{R})}$, from the norm of the successive differences $\Delta\epsilon^N(t)$, provided that M is large enough, and choose N according to the targeted accuracy.

Example 5.2. In the second example, we consider problem (1) with $A(t)$ and $B(t)$ having infinite expansions with coefficients $A_n \sim \text{Beta}(11, 15)$ for $n \geq 0$, $B_0 = 0$, $B_n = 1/n^2$ for $n \geq 1$, while $Y_0 \sim f_{Y_0}(y) = \frac{\sqrt{2}}{\pi(1+y^4)}$ ($-\infty < y < \infty$) and $Y_1 \sim \text{Poisson}(2)$. All these random quantities are assumed to be independent. The power series of $A(t)$ and $B(t)$ converge on $(-1, 1)$ (that is for $r = 1$), so the mean square solution $X(t) = \sum_{n=0}^{\infty} X_n t^n$ given by Theorem 2.1 is defined on $(-1, 1)$. Theorem 3.7 allows approximating $f_{X(t)}(x)$ with $\hat{f}_{X^N(t)}(x)$, $N \geq 0$.

Figure 4 shows graphical representations of $\hat{f}_{X^N(t)}(x)$ for times $t = 0.25, 0.75$ and 0.99 , with orders of truncation $N = 1-5$. The evident regularity of $\hat{f}_{X^N(t)}(x)$ is inherited from the smoothness of the density f_{Y_0} , by Theorem 3.11.




Figure 3: Left: error $E^N(t)$ (see (8)), for different times as indicated. Right: relation between $\log E^N(t)$ and $\log \Delta \epsilon^N(t)$, for $t = 0.5, t = 1$ and $t = 1.5$. Also reported are linear regressions. This figure corresponds to Example 5.1.




Figure 4: Graphical representations of the Monte Carlo estimates $\hat{f}_{X^N(t)}(x)$ at $t = 0.25$ (left), $t = 0.75$ (center) and $t = 0.99$ (right), with orders of truncation N as indicated. This figure corresponds to Example 5.2.

To better assess the convergence, Figure 5 shows the successive differences $\delta \epsilon^N(x, t)$ defined in (6) at the same times as in Figure 4; these differences are decreasing to 0 pointwise as theoretically expected, see Theorem 3.7. As pointwise convergence of densities implies $L^1(\mathbb{R})$ convergence, we report in Table 2 the consecutive norms $\Delta \epsilon^N(t)$ defined by (7). The norms decay, albeit not monotonically; for instance, when $t = 0.25$ the difference is larger for $N = 4$ than in $N = 3$.




Figure 5: Differences in consecutive estimates $\delta \epsilon^N(x, t)$ (see (6)) at $t = 0.25$ (left), $t = 0.75$ (center) and $t = 0.99$ (right), and for orders of truncation as indicated. The plots correspond to Example 5.2.

Figure 6 reports in the left plot the error estimates $\log E^N(t)$ defined in (8). We see that the errors decrease quickly before stagnating because of the sampling error. This example, despite being more complex than the previous one in Example 5.1, in terms of dimensionality, requires smaller orders N , since for $t \in (-1, 1)$ we have $|t - t_0| = |t| < 1$, which implies $|t - t_0|^n \xrightarrow{n \rightarrow \infty} 0$. The

	$N = 1$	$N = 2$	$N = 3$	$N = 4$
$t = 0.25$	0.0215530	0.00607417	0.000600201	0.00167170
$t = 0.75$	0.147952	0.0545970	0.0436801	0.00419704
$t = 0.99$	0.225868	0.0945261	0.127495	0.00985133

Table 2: Norm $\Delta\epsilon^N(t)$ of differences in consecutive estimates (see (7)) for different times t and truncation orders N . This table corresponds to Example 5.2.

right plot of Figure 6 aims at showing the relation between the errors $E^N(t)$ and the successive differences $\Delta\epsilon^N(t)$. Specifically, for the times t shown, a collinearity is found in log-scale through the model (9). In other words, the decay pattern of the consecutive differences characterizes the convergence of the global error as long as the bias error dominates the sampling error.




Figure 6: Left: error $E^N(t)$ in (8), for different times as indicated. Right: relation between $\log E^N(t)$ and $\log \Delta\epsilon^N(t)$, for $t = 0.25, 0.5, 0.75$ and 0.99 . Also reported are linear regressions. This figure corresponds to Example 5.2.

Example 5.3. In this example, we consider the previous degree one polynomial problem, with the following independent distributions: $A_0 = 4$, $A_1 \sim \text{Uniform}(0, 1)$, $B_0 \sim \text{Gamma}(2, 2)|_{[0,4]}$, $B_1 \sim \text{Bernoulli}(0.35)$, $Y_0 \sim \text{Poisson}(2)$ and $Y_1 \sim \text{Normal}(2, 1)$. This example coincides with Example 5.1, except that Y_0 and Y_1 have been interchanged: now Y_0 is discrete, while Y_1 is absolutely continuous. This exchange puts this example in a different theoretical case compared to Example 5.1.

By Theorem 2.1, the unique mean square solution is expressible as a random power series $X(t) = \sum_{n=0}^{\infty} X_n t^n$ that is mean square convergent for all $t \in \mathbb{R}$. According to Remark 3.13, we can approximate the probability density function of $X(t)$, $f_{X(t)}(x)$, for $t \neq 0$.

Figure 7 reports the approximations $\hat{f}_{X^N(t)}(x)$ at times $t = 0.5, 1$ and 1.5 . As N grows, the graphical representations tend to overlap, denoting the convergence of the expansions. The densities are all smooth, as expected from the smoothness of f_{Y_1} , except for $\hat{f}_{X^{N=12}(t=1.5)}(x)$ whose estimate presents noisy features.

The noisy features in $\hat{f}_{X^{N=12}(t=1.5)}(x)$ are due to several reasons. First, there is a computational issue of Mathematica[®] caused by numerical overflow-underflow when too small or too large quantities are involved (for instance $\exp(z)$ for $|z| \gg 1$). Some sample paths of $S_1^{N=12}(t)$ are vanishing near $t = 1.5$, thus making the denominator $S_1^{N=12}(t)$ in the definition of $V_N(t)$ (in (4) by for the role of Y_0 and Y_1 exchanged) very small, with a loss of precision as a result. This is illustrated in Figure 8, where we show some randomly generated sample paths of $S_1^{N=12}(t)$. We also report sample paths for $N = 11$ and $N = 13$ for comparison. Second, and not totally unrelated to the numerical overflow, we have $\mathbb{V}[1/|S_1^N(t = 1.5)|] = \infty$ for $N = 12$ when it remains




Figure 7: Graphical representations of the Monte Carlo estimates $\hat{f}_{X^N(t)}(x)$ at $t = 0.5$ (left), $t = 1$ (center) and $t = 1.5$ (right), with varying orders of truncation N as indicated. This figure corresponds to Example 5.3.

finite for the other values of N shown. As a result, the variance $\sigma_{N=12}^2$ in the Monte Carlo method (see (5)) is unbounded or very large for $N = 12$, while it behaves well for other N , as illustrated in the last panel of Figure 8 (bottom left plot). As a result, for $N = 12$, the convergence of the Monte Carlo procedure is slowed down due to the large or infinite variance, the rate $O(1/\sqrt{M})$ is not obtained (see the discussion from Section 4), and some noisy features plague the estimator.

Luckily, the noise in $\hat{f}_{X^{N=12}(t=1.5)}(x)$ is not present for $N > 12$. In situations where large or infinite variance occurs for some N , one should focus on the truncation orders N for which the approximation $\hat{f}_{X^N(t)}(x)$ behaves nicely, without noise. In this manner, correct approximations to $f_{X(t)}(x)$ are obtained with a feasible number of samples.




Figure 8: Random trajectories of $S_1^N(t)$ for $N = 11, 12$ and 13 . For $N = 12$, observe that some trajectories vanish very close to $t = 1.5$, while for $N \neq 12$ the trajectories remain away from 0. The plot in the bottom right panel shows the corresponding empirical estimates of $\sigma_N^2(x, t = 1.5)$. This figure corresponds to Example 5.3.

Figure 9 (left and center plots) presents the consecutive differences $\delta e^N(x, t)$ given by (6), for times $t = 1$ and 1.5 . These consecutive differences are not monotonically decreasing with N , although a decay pattern towards 0 is perceptible. Further, the impact of the noisy estimate $\hat{f}_{X^{N=12}(t=1.5)}(x)$ is clearly visible in the reported differences. The plots are entirely consistent with the theoretical results and Remark 3.13. In Table 3, we report the corresponding $L^1(\mathbb{R})$ norms

$\Delta\epsilon^N(t)$ (see (7)) as a summary of Figure 9. The last plot of Figure 9 reports the estimate errors $\log E^N(t)$ in (8). Again, the convergence and the sampling error are observed. This example also emphasizes that the Fröbenius method deteriorates for large times, as N needs to increase with t to maintain accurate approximations.




Figure 9: Differences in consecutive estimates $\delta\epsilon^N(x, t)$ at $t = 1$ (left) and $t = 1.5$ (center), with orders of truncation as indicated. The last plot presents the errors $E^N(t)$ (see (8)), for different times as indicated. This figure corresponds to Example 5.3.

	$N = 2$	$N = 3$	$N = 4$	$N = 5$	$N = 6$
$t = 0.5$	0.320171	0.618382	0.333094	0.0893759	0.0291202
	$N = 7$	$N = 8$	$N = 9$	$N = 10$	$N = 11$
$t = 1$	0.308293	0.185148	0.0605758	0.0256469	0.0216692
	$N = 11$	$N = 12$	$N = 13$	$N = 14$	$N = 15$
$t = 1.5$	0.301694	0.155185	0.0677696	0.0375570	0.0202081

Table 3: Norm $\Delta\epsilon^N(t)$ of differences in consecutive estimates (see (7)) for different times t and truncation orders N . This table corresponds to Example 5.3.

Example 5.4. Consider the problem with infinite expansions for A and B , $A_n \sim \text{Beta}(11, 15)$ for $n \geq 0$, $B_0 = 0$, $B_n = 1/n^2$ for $n \geq 1$, $Y_0 \sim \text{Uniform}(-1, 1)$ and $Y_1 \sim \text{Exponential}(2)$. These random inputs are again assumed to be independent. In contrast with Example 5.2, the probability density function of Y_0 has now two discontinuity points at $y_0 = \pm 1$, while Y_1 follows an absolutely continuous law. Hence, Theorem 3.7 cannot be employed here. The mean square analytic solution $X(t) = \sum_{n=0}^{\infty} X_n t^n$ given by Theorem 2.1 is defined on $(-1, 1)$ and we must apply Theorem 3.14 to approximate $f_{X(t)}(x)$ for $t \neq 0$. We compute the approximations at time $t = 0.99$ (near the limit 1), with orders of truncation $N = 1-5$. Figure 10 (left plot) depicts the graphs of $\hat{f}_{X^N(t)}(x)$. Promptly, the successive approximations of the density function tend to superimpose, thus entailing rapid convergence to the target density function $f_{X(t)}(x)$. In contrast to Example 5.2, the non-differentiability of the approximated density functions inherited from f_{Y_0} is evident (here one cannot apply Theorem 3.11). Thereby, our method can capture peaks induced by non-differentiability. This feature is a definite advantage of our method, compared to classical sample paths approximation methods where a kernel density estimation of the density would smear-out the approximation at the non-differentiability points.

A richer analysis of the convergence in this example is provided in the centered plot of Figure 10 and Table 4, which depict consecutive differences $\delta\epsilon^N(x, t)$ (see (6)) and their norms $\Delta\epsilon^N(t)$ (see (7)), respectively. Even though the errors are not decreasing monotonically to 0 (as in the previous Example 5.3), the convergence is evident and follows from Theorem 3.14. Finally, in

Figure 10 (last panel) we also plot the error estimate $\log E^N(t)$ (see (8)), for distinct times. Similar to Example 5.2, the plot shows that this example needs small orders N for all $t \in (-1, 1)$ because of the decay of $|t|^N$.




Figure 10: Graphical representation of the Monte Carlo estimates $\hat{f}_{X^N(t)}(x)$ at $t = 0.99$ (left), with orders of truncation $N = 1-5$. Differences in consecutive estimates $\delta\epsilon^N(x, t)$ (see (6)) at $t = 0.99$ (center), with orders of truncation $N = 1-4$. Error $E^N(t)$ (see (8)), for different times as indicated (right). This figure corresponds to Example 5.4.

	$N = 1$	$N = 2$	$N = 3$	$N = 4$
$t = 0.25$	0.00666810	0.00482414	0.00169730	0.00275401
$t = 0.5$	0.0266447	0.0301975	0.00469989	0.00524461
$t = 0.75$	0.0582143	0.0920184	0.0176119	0.00756466
$t = 0.99$	0.0947395	0.190933	0.0597380	0.0107980

Table 4: Norm $\Delta\epsilon^N(t)$ of differences in consecutive estimates (see (7)) at different times t and for different truncation orders N . This table corresponds to Example 5.4.

Example 5.5. In this final example, we deal with discrete uncertainties, under the setting of Theorem 3.15. Consider again the polynomial problem of Example 5.1, with $A_0 = 4$, $A_1 = 2$, $B_0 = 0$, $B_1 = -1$, and now $Y_0 \sim \text{Bernoulli}(0.4)$ and $Y_1 \sim \text{Uniform}(-1, 1)$, being all independent. By Theorem 2.1, there is a unique mean square solution $X(t) = \sum_{n=0}^{\infty} X_n t^n$ on \mathbb{R} . For each $t \neq 0$, the random variable $X(t)$ is absolutely continuous, due to the absolute continuity of Y_1 . Theorem 3.15, with Y_1 playing the role of Y_0 (see Remark 3.13) allows approximating $f_{X(t)}(x)$ by utilizing the convergence $\lim_{N \rightarrow \infty} \hat{f}_{X^N(t)}(x) = f_{X(t)}(x)$, which holds for almost every $x \in \mathbb{R}$. In this particular example, Algorithm 1 is used with $M = 1,000,000$ iterations, because the deterministic values for $A(t)$ and $B(t)$ make the computational load much less demanding (see the discussion of Section 4). We will thus identify $\hat{f}_{X^N(t)}(x) = f_{X^N, M=1,000,000}^N(x, t)$. For the time $t = 1.5$, the numerical estimates $\hat{f}_{X^N(t)}(x)$ are displayed in Figure 11 (left plot). Observe that the computed density functions are completely different to those of the previous examples: they are discontinuous, in fact step functions, mainly due to the discontinuities in f_{Y_1} . This example highlights the ability of our method to capture discontinuities. The analysis of the convergence is completed with the centered plot from Figure 11 and Table 5, where the consecutive differences $\delta\epsilon^N(x, t)$ (see (6)) and their norms $\Delta\epsilon^N(t)$ (see (7)) are reported, respectively. Table 5 considers times $t = 0.5, 1$ and 1.5 . Finally, the last panel from Figure 11 plots $\log E^N(t)$ (see (8)) as a function of N . The lower bound for the global error is the sampling error, which is smaller than in the previous four examples, owing to the larger number of samples considered. Comparing the last plot from Figure 11 with the corresponding figures from the previous four examples, the

non-monotonic decay of the error is also more pronounced for the three times. The discontinuity in the graph of the target distribution $f_{X(t)}(x)$ can explain the highly non-monotonic decay with the truncation order. Moreover, f_{Y_1} is not Lipschitz continuous, so the exponential convergence rate discussed in Remark 3.10 is not applicable in the present example. Finally, as for the other examples, the truncation order needed to reduce the error to the sampling contribution increases as we move away from the origin $t_0 = 0$. This behavior may pose severe challenges for large times t .




Figure 11: In the left plot, graphical representations of the Monte Carlo estimates $\hat{f}_{X^N(t)}(x)$ at $t = 1.5$, with orders of truncation N as indicated. In the center plot, differences in consecutive estimates $\delta\epsilon^N(x, t)$ (see (6)) at $t = 1.5$. In the last plot, error $E^N(t)$ (see (8)), for different times as indicated. This figure corresponds to Example 5.5.

	$N = 2$	$N = 3$	$N = 4$	$N = 5$	$N = 6$
$t = 0.5$	0.833333	0.678571	0.140704	0.0280760	0.0190810
	$N = 7$	$N = 8$	$N = 9$	$N = 10$	$N = 11$
$t = 1$	0.785425	0.0767401	0.187360	0.0665995	0.00866170
	$N = 15$	$N = 16$	$N = 17$	$N = 18$	$N = 19$
$t = 1.5$	0.331941	0.0569325	0.0732120	0.0390644	0.00346696

Table 5: Norm $\Delta\epsilon^N(t)$ of differences in consecutive estimates (see (7)) at different times t and truncation orders N . This table corresponds to Example 5.5.

6. Conclusions and perspectives

In this paper, we address the analysis of the random non-autonomous second order linear differential equation. When the data $A(t)$ and $B(t)$ are given by random power series on $(t_0 - r, t_0 + r)$ in $L^\infty(\Omega)$, say $A(t) = \sum_{n=0}^{\infty} A_n(t - t_0)^n$ and $B(t) = \sum_{n=0}^{\infty} B_n(t - t_0)^n$, and the initial conditions Y_0 and Y_1 belong to $L^2(\Omega)$, it is possible to construct a random power series solution $X(t) = \sum_{n=0}^{\infty} X_n(t - t_0)^n$ on $(t_0 - r, t_0 + r)$ in the mean square sense, whose coefficients satisfy a random difference equation. This approach is the generalization of the Fröbenius method to the random setting. The convergence rate of the power series of $X(t)$ is exponential for each time t , but not uniformly on the whole time domain $(t_0 - r, t_0 + r)$. For a fixed tolerance on the mean square error of $X(t)$, the order of truncation of the power series needs to be increased, in general, when $|t - t_0|$ grows.

We proved the existence of solution $X(t)$ when $A(t)$ and $B(t)$ are bounded. In the unbounded case, we present in this paper two counterexamples of existence. In practice, the hypotheses can be fulfilled truncating unbounded random coefficients.

The probability density function of $X(t)$ can be expressed as the expectation of a random process $Z(x, t)$, $f_{X(t)}(x) = \mathbb{E}[Z(x, t)]$, using the law of total probability. A closed-form expression for Z is derived in terms of the fundamental set by exploiting the linearity of the system. However, to compute this expectation, one needs to perform a dimension reduction of the problem, by truncating the series of the fundamental set used to express the solution $X(t)$. Denoting $X^N(t)$, $N \geq 0$, the truncation of $X(t)$, we show that $f_{X^N(t)}(x)$ converges to $f_{X(t)}(x)$ pointwise as $N \rightarrow \infty$ under certain conditions (regarding Nemytskii operators); in some cases, an exponential convergence may be achieved for each t and x . The pointwise convergence also implies convergence in $L^p(\mathbb{R})$, $1 \leq p < \infty$. In particular, the convergence in $L^1(\mathbb{R})$ is equivalent to the convergence in the total variation and the Hellinger distances, which are instances of f -divergences.

From a numerical standpoint, the expectation defining $f_{X^N(t)}(x) = \mathbb{E}[Z^N(x, t)]$ is computable *via* a classical Monte Carlo strategy. We propose an algorithm for that purpose, which estimates symbolically $f_{X^N(t)}(x)$. This algorithm is implemented in the software *Mathematica*[®], and it can be used to compute pointwise approximations of the density function $f_{X(t)}(x)$. One key feature of the algorithm is that it handles discontinuity and non-differentiability points of $f_{X(t)}(x)$ appropriately, without smoothing them out.

To the best of our knowledge, this paper is the first one to provide such analysis of random second order linear differential equations. However, we point out certain limitations of our methodology, which constitute potential avenues for future developments.

To start with, despite the exponential convergence rate, the approximations substantiated on the Fröbenius method may deteriorate for large $|t - t_0|$. This fact is inherent to Taylor series-based methods and also plagues other types of stochastic computations, such as PC expansions. Following [47], using random time-transformations may help to improve the convergence of the Fröbenius method and mitigate this issue.

Another point requiring a more in-depth analysis is the ignorance of the specific values of δ and μ . In particular, we showed that if the truncated processes from the fundamental set vanish for some trajectories near the time t of interest, the numerical estimate of the density becomes very noisy (see Example 5.3). This effect is due to the variance of $Z^N(x, t)$ that may be very large or infinite, with a severely deteriorated Monte Carlo convergence in these situations. We are currently exploring different strategies to sort out this issue, such as the path-wise selection of the variable (Y_0 or Y_1) used in the expression of $Z^N(x, t)$, in order to control its variance.

Efforts to weaken or modify the theoretical hypotheses and enlarge the applicability of our method shall also be carried out. As an example, the extension of the method to the case of Y_0 and Y_1 not absolutely continuous would also present a valuable achievement. Similarly, an extension of the present methodology to linear systems of second order random differential equations may be of great interest, while the application to other stochastic models of our expertise on random expansions and density approximations could be interesting.

At the computational level, the Monte Carlo estimation of $f_{X^N(t)}(x)$ introduces a statistical error since, in numerical computations, we are restricted to a finite number M of realizations. Therefore, an error of order $1/\sqrt{M}$ is unavoidable, even for $N \approx \infty$. The results presented in the paper have highlighted the crucial importance of bias and sampling errors. In the future, it would be beneficial to rely on improved sampling strategies, such as multilevel Monte Carlo [48, 49], to balance the bias and sampling errors, while reducing the computational cost of the Monte Carlo estimates of the density. This topic is the focus of our current efforts.

Acknowledgements

This work is supported by the Spanish “Ministerio de Economía y Competitividad” grant MTM2017–89664–P. M.J. acknowledges the doctorate scholarship granted by PAID, as well as “Ayudas para movilidad de estudiantes de doctorado de la Universitat Politècnica de València para estancias en 2019”, for financing his research stay at CMAP. J.C. acknowledges “Fundació Ferran Sunyer i Balaguer”, “Institut d’Estudis Catalans” and the award from “Borses Ferran Sunyer i Balaguer 2019” for funding her research stay at CMAP. All authors are also grateful to Inria (Centre de Saclay, DeFi Team), which hosted M.J. and J.C. during their research stays at École Polytechnique.

References

- [1] T.T. Soong, *Random Differential Equations in Science and Engineering*; Academic Press: New York, 1973.
- [2] T. Neckel, F. Rupp. *Random Differential Equations in Scientific Computing*. München, Germany: Walter de Gruyter, 2013.
- [3] J.L. Strand, *Random ordinary differential equations*. *J. Differ. Equ.* 1970, 7, 538–553.
- [4] Villafuerte, L., Braumann, C. A., Cortés, J. C., Jódar, L. (2010). Random differential operational calculus: theory and applications. *Computers & Mathematics with Applications*, 59(1), 115–125.
- [5] R. C. Smith. *Uncertainty Quantification. Theory, Implementation, and Application*. SIAM Computational Science & Engineering, 2014.
- [6] G. Fishman. *Monte Carlo: Concepts, Algorithms, and Applications*. Springer Science & Business Media, 2013.
- [7] R. Ghanem, P.D. Spanos. *Stochastic Finite Elements: A Spectral Approach*. Springer-Verlag, New York, 1991.
- [8] D. Xiu. *Numerical Methods for Stochastic Computations. A Spectral Method Approach*. Cambridge Texts in Applied Mathematics, Princeton, University Press, New York, 2010.
- [9] Le Maître O., Knio O. M. (2010). *Spectral Methods for Uncertainty Quantification: With Applications to Computational Fluid Dynamics*. Springer Science & Business Media.
- [10] Liu W.K., Belytschko T., Mani A. (1986). Probabilistic finite elements for nonlinear structural dynamics. *Computer Methods in Applied Mechanics and Engineering*, 56(1), 61–81.
- [11] J. Licea, L. Villafuerte, B. M. Chen-Charpentier. Analytic and numerical solutions of a Riccati differential equation with random coefficients. *Journal of Computational and Applied Mathematics*, 309(1) (2013), 208–219.
- [12] M. A. El-Tawil. The approximate solutions of some stochastic differential equations using transformations. *Applied Mathematics and Computation*, 164(1) (2005), 167–178.
- [13] Dorini, F., Cunha, M.: Statistical moments of the random linear transport equation. *J. Comput. Phys.* 227(19), 8541–8550 (2008).
- [14] Hussein, A., Selim, M.M.: A developed solution of the stochastic Milne problem using probabilistic transformations. *Appl. Math. Comput.* 216(10), 2910–2919 (2009).
- [15] J. Calatayud, J.-C. Cortés, M. Jornet. The damped pendulum random differential equation: A comprehensive stochastic analysis via the computation of the probability density function. *Physica A: Statistical Mechanics and Applications*, 512, 261–279 (2018).
- [16] Nouri, K., Ranjbar, H., Mean square convergence of the numerical solution of random differential equations. *Mediterr. J. Math.* 12(3), 1123–1140 (2015).
- [17] Cortés, J.-C.; Jódar, L.; Camacho, J.; Villafuerte, L. Random Airy type differential equations: Mean square exact and numerical solutions. *Comput. Math. Appl.* 2010, 60, 1237–1244.
- [18] Calbo, G., Cortés, J.C., Jódar, L.: Random Hermite differential equations: mean square power series solutions and statistical properties. *Appl. Math. Comput.* 218(7), 3654–3666 (2011).
- [19] J. Calatayud, J.-C. Cortés, M. Jornet. Improving the approximation of the first and second order statistics of the response stochastic process to the random Legendre differential equation. *Mediterranean Journal of Mathematics*, 16:68 (2019).
- [20] J. Calatayud, J.-C. Cortés, M. Jornet, L. Villafuerte. Random non-autonomous second order linear differential equations: mean square analytic solutions and their statistical properties. *Advances in Difference Equations*, 2018:392, 1–29 (2018).
- [21] J. Calatayud, J.-C. Cortés, M. Jornet. Some Notes to Extend the Study on Random Non-Autonomous Second Order Linear Differential Equations Appearing in Mathematical Modeling. *Mathematical and Computational Applications*, 23(4), 76–89 (2018).

- [22] J. Calatayud, J.-C. Cortés, M. Jornet. Computational uncertainty quantification for random non-autonomous second order linear differential equations via adapted gPC: a comparative case study with random Fröbenius method and Monte Carlo simulation. *Open Mathematics*, 16, 1651–1666 (2018).
- [23] Golmankhaneh, A.A.; Porghoveh, N.A.; Baleanu, D. Mean square solutions of second-order random differential equations by using homotopy analysis method. *Rom. Rep. Phys.* 2013, 65, 350–362.
- [24] Khudair, A.R.; Ameen, A.A.; Khalaf, S.L. Mean square solutions of second-order random differential equations by using variational iteration method. *Appl. Math. Sci.* 2011, 5, 2505–2519.
- [25] Khudair, A.R.; Ameen, A.A.; Khalaf, S.L. Mean square solutions of second-order random differential equations by using Adomian decomposition method. *Appl. Math. Sci.* 2011, 5, 252–2535.
- [26] A. K. Khudair, S. A. M. Haddad, S. L. Khalaf. Mean square solutions of second-order random differential equations by using the differential transformation method. *Open Journal of Applied Sciences* 6 (2016), 287–297.
- [27] Villafuerte, L.; Chen-Charpentier, B.-M. A random differential transform method: Theory and applications. *Appl. Math. Lett.* 2012, 25, 1490–1494.
- [28] Casabán, M.C., Cortés, J.C., Romero, J.V., Roselló, M.D.: Solving random homogeneous linear second-order differential equations: a full probabilistic description. *Mediterr. J. Math.* 13(6), 3817–3836 (2016).
- [29] Cortés, J. C., Navarro-Quiles, A., Romero, J. V., Roselló, M. D. (2018). Solving second-order linear differential equations with random analytic coefficients about ordinary points: A full probabilistic solution by the first probability density function. *Applied Mathematics and Computation*, 331, 33–45.
- [30] Gerritsma, M., Van der Steen, J. B., Vos, P., Karniadakis, G. (2010). Time-dependent generalized polynomial chaos. *Journal of Computational Physics*, 229(22), 8333–8363.
- [31] J.L. Doob, *Stochastic Processes*, John Wiley & Sons, 1953.
- [32] R.B. Ash. *Real Analysis and Probability*. Academic Press, New York, 1972.
- [33] D. Nualart, M. Sanz. *Curs de Probabilitats (in Catalan)*. PPU, Barcelona, 1990.
- [34] Calatayud, J.; Cortés, J.C.; Jornet, M. Uncertainty quantification for random parabolic equations with non-homogeneous boundary conditions on a bounded domain via the approximation of the probability density function. *Math. Method Appl. Sci.* 2018.
- [35] J. Calatayud, B.M. Chen-Charpentier, J.-C. Cortés, M. Jornet. Combining polynomial chaos expansions and the random variable transformation technique to approximate the density function of stochastic problems, including some epidemiological models. *Symmetry*, 11(1), 43 (2019).
- [36] D. Williams. *Probability with Martingales*. New York: Cambridge University Press, 1991.
- [37] H. Scheffé. A Useful Convergence Theorem for Probability Distributions. *The Annals of Mathematical Statistics*, 18(3): 434–438, 1947.
- [38] L. Tenorio. *An Introduction to Data Analysis and Uncertainty Quantification for Inverse Problems*. Vol. 3, SIAM, 2017.
- [39] E. Hellinger, Neue Begründung der Theorie quadratischer Formen von unendlichvielen Veränderlichen, *Journal für die reine und angewandte Mathematik*, 136: 210–271 (1909).
- [40] A. Ambrosetti, G. Prodi, *A Primer of Nonlinear Analysis*. Cambridge Studies in Advanced Mathematics, 1993.
- [41] M.M. Vainberg, *Variational Methods for the Study of Nonlinear Operators*, Holden-Day, 1964.
- [42] Haim Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, 2011.
- [43] A.W. van der Vaart, *Asymptotic Statistics*, Cambridge University Press, 1998.
- [44] Klenke, A. (2013). *Probability Theory: A Comprehensive Course*. 2nd ed, Springer Science & Business Media.
- [45] K.O. Geddes, S.R. Czapor, G. Labahn (1992). *Algorithms for Computer Algebra*.
- [46] Wolfram Research, Inc, 2017. *Mathematica*, Version 11.2; Wolfram Research, Inc.: Champaign, IL, USA.
- [47] O.P Le Maître, L. Mathelin, O.M. Knio and Y.M. Hussaini (2010). Asynchronous time integration for polynomial Chaos expansions of uncertain periodic dynamics. *Discrete and Continuous Dynamical Systems*, 28(1), 199–226.
- [48] Giles, M. B. (2008). Multilevel Monte Carlo path simulation. *Operations Research*, 56(3), 607–617.
- [49] Giles, M. B. (2015). Multilevel Monte Carlo methods. *Acta Numerica*, 24, 259–328.